arXiv:math/9805070v1 [math.DG] 14 May 1998

L^2-torsion of hyperbolic manifolds

Eckehard Hess Thomas Schick

to appear in Manuscripta Mathematica

Eckehard Hess
Fachbereich Mathematik
Universität Mainz

55099 Mainz, Germany
hess@topologie.mathematik.uni-mainz.de

Thomas Schick
Fachbereich Mathematik
Universität Münster
Einsteinstr. 62
48149 Münster, Germany
thomas.schick@math.uni-muenster.de

The L^2-torsion is an invariant defined for compact L^2-acyclic manifolds of determinant class, for example odd dimensional hyperbolic manifolds. It was introduced by John Lott [Lot92] and Varghese Mathai [Mat92] and computed for hyperbolic manifolds in low dimensions. Our definition of the L^2-torsion coincides with that of John Lott, which is twice the logarithm of that of Varghese Mathai.

In this paper you will find a proof of the fact that the L^2-torsion of hyperbolic manifolds of arbitrary odd dimension does not vanish. This was conjectured by John Lott in [Lot92, p.484, Proposition 16 infra]. Some concrete values are computed and an estimate of their growth with the dimension is given. The values we compute for dimensions 5 and 7 differ from those published in [Lot92, Proposition 16]. The result has been independently achieved by both authors and will be part of the dissertation of Eckehard Hess at the university of Mainz. For an introduction into L^2-theory see [Lü97].

We are indebted to Prof. Dr. Lück, Münster, for permanent support and encouragement.

1. Definition. Following [Lot92, p.482] we define the analytic L^2-torsion of an L^2-acyclic Riemannian $(d = 2n+1)$-dimensional manifold of determinant class by

$$\mathrm{Tor}_{(2)}(M) = 2\sum_{j=0}^{n}(-1)^{j+1}\operatorname{logdet}_G(\triangle_j|)$$

Here G is the fundamental group of M, $\triangle_j|$ is the Laplacian restricted to coclosed forms on the universal covering $\tilde{M}$ and the logarithm of the G-determinant is computed from the local trace of the heat kernel as follows

$$\begin{aligned}\operatorname{logdet}_G(\triangle_j|) \;=\; & \int_F \left\{ \frac{d}{ds}\bigg|_{s=0} \left[\frac{1}{\Gamma(s)}\int_0^1 t^{s-1}\mathrm{tr}_{\mathbb{C}} e^{-t\triangle_j|}(x,x)dt \right] \right. \\ & \left. + \int_1^\infty t^{-1}\mathrm{tr}_{\mathbb{C}} e^{-t\triangle_j|}(x,x)dt \right\} dx\end{aligned}$$

Here F is a fundamental domain of M in $\tilde{M}$, the first integral exists for s sufficiently large and one has to take the meromorphic extension at 0. M being of determinant class ensures the second integral to converge.

2. Theorem. *There is a constant $\alpha_d > 0$, such that for every $(d = 2n+1)$-dimensional closed hyperbolic manifold*

$$\operatorname{Tor}_{(2)}(M) = (-1)^n \alpha_d \operatorname{Vol}(M)$$

The values for α_d have been computed as follows. Although exact values were computed for $d \leq 251$, we will give only exact numbers for $d \leq 11$ and approximate numbers for $d \leq 39$:

d	α_d	$\approx \alpha_d$
3	$\frac{1}{3\pi}$	$0,106103$
5	$\frac{62}{45\pi^2}$	$0,139598$
7	$\frac{221}{35\pi^3}$	$0,203645$
9	$\frac{32204}{945\pi^4}$	$0,349847$
11	$\frac{1339661}{6237\pi^5}$	$0,701891$

d	$\approx \alpha_d$
13	$1,61885$
15	$4,22925$
17	$12,3578$
19	$39,9606$
21	$141,729$
23	$547,188$

25	$2284,87$
27	$10261,5$
29	49326
33	252701
35	$1,37458 * 10^6$
37	$7,91236 * 10^6$
39	$4,80523 * 10^7$

3. Lemma. *Let $\triangle_j| = \delta_j d_j|_{ker(\delta_{j+1})}$ be the Laplacian, restricted to coclosed L^2-forms on the $(d = 2n+1)$-dimensional hyperbolic space H^d. Then for every closed d-dimensional hyperbolic manifold M with fundamental group G and $j \neq n$*

$$\operatorname{logdet}_G(\triangle_j|) = \operatorname{Vol}(M) C \binom{2n}{j} \sum_{k=0}^{n} K^n_{k,j} (-1)^{k+1} \tfrac{2\pi}{2k+1} (n-j)^{2k+1}$$

$$\textit{with } C = \frac{(4\pi)^{-(n+\frac{1}{2})}}{\Gamma(n+\frac{1}{2})} \qquad \textit{and} \qquad a = n-j$$

Here $K^n_{k,j}$ is the coefficient of ν^{2k} in the polynomial

$$P(\nu) := \frac{\prod_{i=0}^{n}(\nu^2 + i^2)}{\nu^2 + (n-j)^2} \qquad (*)$$

Note that $P(\nu)$ indeed is a polynomial rather than a rational function, as $|a| \in \{1...n\}$. In addition

$$\operatorname{logdet}_G(\triangle_n|) = 0$$

Proof. Following [Lot92, prop. 15] the local trace of the heat kernel of $\triangle_j|$ is

$$\operatorname{tr}_{\mathbb{C}} e^{-t\triangle_j|}(x,x) = C \binom{2n}{j} \int_{-\infty}^{\infty} e^{-t(\nu^2+a^2)} \frac{\prod_{k=0}^{n}(\nu^2+k^2)}{\nu^2+a^2} d\nu$$

According to the above remark let

$$\frac{\prod_{k=0}^{n}(\nu^2+k^2)}{\nu^2+a^2} = \sum_{k=0}^{n} K^n_{k,j} \nu^{2k}$$

Evaluation of the above integral yields

$$\mathrm{tr}_{\mathbb{C}} e^{-t\triangle_j}(x,x) = C\left(\begin{array}{c} 2n \\ j \end{array}\right) \sum_{k=0}^{n} K_{k,j}^{n} e^{-ta^2} t^{-k-\frac{1}{2}} \Gamma\Big(k+\frac{1}{2}\Big)$$

Now we have to compute

$$\begin{aligned} L_j &:= \frac{\mathrm{logdet}(\triangle_j|)}{\mathrm{Vol}(M)} \\ &= \frac{d}{ds}\bigg|_{s=0} \left[\frac{1}{\Gamma(s)} \int_0^1 t^{s-1} \mathrm{tr}_{\mathbb{C}} e^{-t\triangle_j}(x,x) dt\right] + \int_1^{\infty} t^{-1} \mathrm{tr}_{\mathbb{C}} e^{-t\triangle_j}(x,x) dt \end{aligned}$$

John Lott showed in [Lot92, Lemma 13, p.481]

$$L_n = 0$$

For $j \neq n$, that is $a = n - j \neq 0$

$$L_j = C\left(\begin{array}{c} 2n \\ j \end{array}\right) \sum_{k=0}^{n} K_{k,j}^{n} \Gamma\Big(k+\frac{1}{2}\Big) \frac{d}{ds}\bigg|_{s=0} \left(\frac{1}{\Gamma(s)} \underbrace{\int_0^{\infty} e^{-ta^2} t^{s-k-\frac{3}{2}} dt}_{:=J}\right)$$

J exists for s sufficiently large. Its meromorphic extension leads to

$$\begin{aligned} L_j &= C\left(\begin{array}{c} 2n \\ j \end{array}\right) \sum_{k=0}^{n} K_{k,j}^{n} \Gamma\Big(k+\frac{1}{2}\Big) \Gamma\Big(-k-\frac{1}{2}\Big) a^{2k+1} \\ &= C\left(\begin{array}{c} 2n \\ j \end{array}\right) \sum_{k=0}^{n} K_{k,j}^{n} (-1)^{k+1} \tfrac{2\pi}{2k+1} a^{2k+1} \end{aligned}$$

□

4. Corollary. *For any closed hyperbolic manifold of dimension* $d = 2n+1$ *we have by Definition 1*

$$\frac{\mathrm{Tor}_{(2)}(M)}{\mathrm{Vol}(M)} = 2 \sum_{j=0}^{n-1} (-1)^{j+1} C\left(\begin{array}{c} 2n \\ j \end{array}\right) \sum_{k=0}^{n} K_{k,j}^{n} (-1)^{k+1} \tfrac{2\pi}{2k+1} (n-j)^{2k+1}$$

The numerical values were computed using this fomula and Mathematica.

5. Lemma. *Let* M *be a closed* $(d = 2n+1)$*-dimensional manifold. Then*

$$(-1)^{j+1} \mathrm{logdet}_G(\triangle_j|) = (-1)^n |\mathrm{logdet}_G(\triangle_j|)|$$

with

$$|\mathrm{logdet}_G(\triangle_j|)| > 0 \textit{ for } j \neq n$$

In particular

$$(-1)^n \,\mathrm{Tor}_{(2)}(M) > 0$$

Proof. Let $j \neq n$. Then we have

$$\begin{aligned} L_j &= -2\pi C \binom{2n}{j} \sum_{k=0}^{n} K_{k,j}^n (-1)^k \tfrac{1}{2k+1} a^{2k+1} \\ &= -2\pi C \binom{2n}{j} \int_0^a \sum_{k=0}^{n} K_{k,j}^n (ix)^{2k} dx \end{aligned}$$

Using the definition $(*)$ of the coefficients $K_{k,j}^n$ in Lemma 3 one gets

$$L_j = -2\pi C \binom{2n}{j} \int_0^a \frac{\prod_{k=0}^n (k^2 - x^2)}{a^2 - x^2} dx$$

One has

$$\begin{aligned} &\int_0^a \frac{\prod_{k=0}^n (k^2 - x^2)}{a^2 - x^2} dx \\ &= (-1)^{n+1} \int_0^a \frac{x}{(a+x)(a-x)} \prod_{k=-n}^{n} (x+k) dx \\ &= (-1)^{n+1} \sum_{r=0}^{a-1} \int_0^1 f_r(t) dt \end{aligned}$$

where for $t \in]0,1[\ ,\ r \in \{0,..,a-1\}$ we define

$$\begin{aligned} f_r(t) &= \underbrace{\frac{t+r}{(a+t+r)(a-t-r)}}_{>0} \prod_{k=-n+r}^{n+r} \underbrace{(t+k)}_{\substack{<0 \text{ for } k<0 \\ >0 \text{ otherwise}}} \\ &= (-1)^{n-r} |f_r(t)| \end{aligned}$$

For $t \in]0,1[$ and $0 \leq r < r+1 \leq a-1$ one computes

$$\left| \frac{f_{r+1}(t)}{f_r(t)} \right| > 1$$

Hence

$$\int_0^1 |f_{r+1}(t)| dt \geq \int_0^1 |f_r(t)| dt$$

Now the sum

$$L_j = -2\pi C \binom{2n}{j} \sum_{r=0}^{a-1} (-1)^{r+1} \int_0^1 |f_r(t)| dt$$

is an alternating sum and the absolute values of the summands are stricly increasing. So it is not 0 and the sign is that of the last summand. One concludes

$$\operatorname{logdet}_G(\triangle_j|) = (-1)^{n-j-1} |\operatorname{logdet}_G(\triangle_j|)|$$

This also finishes the proof of Theorem 2. $\square$

6. Proposition. *The constants* α_d *of Theorem 2 strictly increase and*

$$\alpha_{2n+1} \geq \frac{n}{2\pi}\alpha_{2n-1}$$

In particular

$$\alpha_{2n+1} \geq \frac{2}{3}\frac{n!}{(2\pi)^n}$$

Proof. An elementary computation shows

$$\left|\frac{f_{a-1}(t)}{f_{a-2}(t)}\right| \geq 2$$

Now one has

$$\left|\int_0^a \frac{\prod_{k=0}^n (k^2-x^2)}{a^2-x^2}dx\right| \;\geq\; \frac{1}{2}\int_{a-1}^a \left|\frac{\prod_{k=0}^n (k^2-x^2)}{a^2-x^2}\right| dx$$

and

$$\begin{aligned}
\alpha_{2n+1} &\geq 2\pi\sum_{j=0}^{n-1}\frac{(4\pi)^{-(n+\frac{1}{2})}}{\Gamma\Big(n+\frac{1}{2}\Big)}\begin{pmatrix}2n\\ j\end{pmatrix}\int_{n-j-1}^{n-j}\left|\frac{\prod_{k=0}^{n-1}(k^2-x^2)}{(n-j)^2-x^2}\right|\underbrace{(n^2-x^2)}_{\geq(2n-j)j}dx \\
&\geq 2\pi\sum_{j=1}^{n-1}\frac{(4\pi)^{-(n+\frac{1}{2})}}{\Gamma\Big(n+\frac{1}{2}\Big)}2n(2n-1)\begin{pmatrix}2n-2\\ j-1\end{pmatrix}\int_{n-j-1}^{n-j}\left|\frac{\prod_{k=0}^{n-1}(k^2-x^2)}{(n-j)^2-x^2}\right|dx \\
&= 4\pi\sum_{j=1}^{n-1}\frac{2n}{4\pi}\frac{(4\pi)^{-(n-1+\frac{1}{2})}}{\Gamma\Big(n-1+\frac{1}{2}\Big)}\begin{pmatrix}2n-2\\ j-1\end{pmatrix}\int_{n-j-1}^{n-j}\left|\frac{\prod_{k=0}^{n-1}(k^2-x^2)}{(n-j)^2-x^2}\right|dx \\
&\geq \frac{n}{2\pi}\,4\pi\sum_{l=0}^{n-2}\frac{(4\pi)^{-(n-1+\frac{1}{2})}}{\Gamma\Big(n-1+\frac{1}{2}\Big)}\begin{pmatrix}2(n-1)\\ l\end{pmatrix}\int_{n-1-l-1}^{n-1-l}\left|\frac{\prod_{k=0}^{n-1}(k^2-x^2)}{(n-1-l)^2-x^2}\right|dx \\
&\geq \frac{n}{2\pi}\,4\pi\sum_{l=0}^{n-2}\frac{(4\pi)^{-(n-1+\frac{1}{2})}}{\Gamma\Big(n-1+\frac{1}{2}\Big)}\begin{pmatrix}2(n-1)\\ l\end{pmatrix}\left|\int_0^{n-1-l}\frac{\prod_{k=0}^{n-1}(k^2-x^2)}{(n-1-l)^2-x^2}dx\right| \\
&= \underbrace{\frac{n}{2\pi}}_{\geq 1 \text{ for } n\geq 7}\quad\alpha_{2n-1}
\end{aligned}$$

For $n \leq 7$ the growth follows from the table. ☐

References

[Lü97] W. Lück. L^2-invariants of regular coverings of compact manifolds and *CW*-complexes. In R.B. Daverman, and R.J. Sher, editors, *Handbook of Geometry*. Elsevier, to appear.

[Lot92] John Lott. Heat kernels on covering spaces and topological invariants. *Journal of Differential Geometry*, 35:471–510, 1992.

[Mat92] Varghese Mathai. L^2-analytic torsion. *Journal of Functional Analysis*, 107(2):369–386, 1992.